

# A COMPUTATIONAL STUDY OF MECHANISTIC PATHWAYS FOR THE DESIGN OF A GREENER PROCESS IN THE PRODUCTION OF QUINOLINE-2-(1H)-ONE AND ITS DERIVATIVES

**Blessing F. Masora**<sup>1</sup>, Krishna K. Govender<sup>1\*</sup>, Penny P. Govender<sup>1</sup>, Ephraim F. Marondedze<sup>1</sup>

<sup>1</sup>*Department of Chemical Sciences, University of Johannesburg, Johannesburg, South Africa.*

\*Corresponding Author Email: krishnag@uj.ac.za

## ABSTRACT

Quinolinones are heterocyclic aromatic compounds with various pharmacological activities. Their importance has brought about the need for green synthetic methods [1]. Exploring greener pathways using an experimental trial and error approach is costly and hazardous hence computational investigations of the existing mechanistic pathways are an alternative [2]. This study focused on the computational study of a fast, base-free aqueous synthesis of quinolin-2(1H)-one. Molecular simulations were done using DFT [3], MP2, and CCSD(T) [4] to obtain reaction energy profiles of quinoline-2(1H)-one in gas, water, and dichloromethane with the aid of Gaussian 16 revision C01[5]. The computational simulations were conducted at the Centre for High Performance Computing (CHPC). The jobs were submitted for execution on a single node with 24 cores and a specified memory of 10GB. The scheduled wall times utilized varied between 10-96 hrs for optimisation and normal mode analysis, which were executed as two separate simulations for each species investigated. The profiles for the two-step reaction helped to elucidate the energy barriers for the chemical reaction. The barrier for the reaction in water and dichloromethane in the first step was 22.44 and 20.58 kcal/mol, respectively, while for the second step 27.24 and 28.35 kcal/mol was obtained. HOMO-LUMO analysis and quantum theory of atoms in molecules (QTAIM) calculations were explored to provide further insight into the reaction. The energy gap for the final product was 0.249 eV indicating a relatively stable molecule which could possibly participate in other reactions. The stability was supported by average rho and Laplacian of 0.253 and -0.558, respectively, via QTAIM analysis.

**Keywords:** quinoline-2(1H)-one, green, CHPC, computational, pathway, simulation.

## References

1. Julie B. Zimmerman, Paul T. Anastas, Hanno C. Erythropel, Walter Leitner. Designing for a green chemistry future. *Science*, 2020; 367 (6476): 397.
2. Yang, Jacob N. Sanders, Wei Li, Peiyuan Yu, Hongbo Wang, Zhijun Tang, Wen Liu, and K. N. Houk. *Journal of the American Chemical Society* 2020 142 (47), 20232-20239
3. Zhang Y, Liu J, Ghoshal S, Moores A. Density Functional Theory Calculations Decipher Complex Reaction Pathways of 6:2 Fluorotelomer Sulfonate to Perfluoroalkyl Carboxylates Initiated by Hydroxyl Radical. *Environ Sci Technol.* 2021;55(24):16655-16664
4. M. Loipersberger, L. W. Bertels, J. Lee, and M. Head-Gordon, "Exploring the Limits of Second- and Third-Order Møller-Plesset Perturbation Theory for Non-Covalent Interactions: Revisiting MP2.5 and Assessing the Importance of Regularization and Reference Orbitals," *Journal of chemical theory and computation*, vol. 17, no. 9, p. 5582, Sep. 2021, doi: 10.1021/ACS.JCTC.1C00469.
5. M. J. Frisch et al., "Gaussian 16 Revision C.01," Gaussian, Inc., Wallingford CT, 2016.